

Efficon®

Insecticide - Powered by Axalion® Active

INSECTICIDES


| COMPATIBILITY WITH EFFICON® | | | |
|--|---|---|---|
| Abamectin 1.9% EC | ● | Thiamethoxam 12.6% + Lambda cyhalothrin 9.5% ZC | ● |
| Imidacloprid 70% WG | ● | Novaluron 5.25% + Emamectin benzoate 0.9% SC | ● |
| Chlorantraniliprole 18.5% SC | ● | Ethion 50% EC | ● |
| Broflanilide 30% SC | ● | Chlорfenapyr 10% SC | ● |
| Flubendiamide 48% SC | ● | Cyenopyrafen 30% SC | ● |
| Pyriproxyfen 10% EC | ● | Spirotetramet 11.01% + Imidacloprid 11.01% SC | ● |
| Emamectin Benzoate | ● | Spiromesifen 22.9% SC | ● |
| Fipronil 5% SC | ● | Flubendiamide 19.92% + Thiacloprid 19.92% SC | ● |
| Spinosad 48% SC | ● | Lambda-Cyhalothrin 5% EC | ● |
| Flonicamid 50% WG | ● | Imidacloprid 40% + Fipronil 40% w/w WG | ● |
| Fenazaquin 10% EC | ● | Afidopyropen 5% DC | ● |
| Carbosulfan 25% EC | ● | Fenpyroximate 5% EC | ● |
| Propargite 57% EC | ● | Dimethoate 30% EC | ● |
| Thiodicarb 75% WP | ● | Novaluron 5.25% + Emamectin benzoate 0.9% SC | ● |
| Emamectin Benzoate 5% + Lufenuron 40% WG | ● | Chloropyrifos 50% EC | ● |
| Cyantraniliprole 10.26% OD | ● | Monocrotophos 36% SL | ● |
| Etoxazole 10% SC | ● | Profenofos 50% EC | ● |
| Thiacloprid 24% SC | ● | Cypermethrin 5% EC | ● |
| Thiamethoxam 25% WG | ● | | |
| Spinetoram 11.7% SC | ● | Carbosulfan 25% EC | ● |
| Quinolphos 20% + Cypermethrin 3% E.C. | ● | Chloropyrifos 50% + Cypermethrin 5% EC | ● |
| Chlorfluazuron 5.4% EC | ● | Chlorantraniliprole 9.3% + Lambda-Cyhalothrin 4.6% ZC | ● |
| Acephate 50% + Imidacloprid 5% WP | ● | Profenophos 50% + Cypermethrin 5% EC | ● |

FUNGICIDES

| COMPATIBILITY WITH EFFICON® | | | |
|--|---|--|---|
| Azoxystrobin 18.2% + Difenoconazole 11.4% SC | ● | Myclobutanil 10% WP | ● |
| Azoxystrobin 20% + Cyproconazole 8% SC | ● | Copper Hydroxide 53.8% DF | ● |
| Carbendazim 50% WP | ● | Fluopyram 17.7% + Tebuconazole 17.7% SC | ● |
| Metiram 55% + Pyraclostrobin 5% WG | ● | Flusilazole 12.5% + Carbendazim 25% SE | ● |
| Copper Oxychloride 50% WP | ● | Boscalid 25.2% + Pyraclostrobin 12.8% WG | ● |
| Kasugamycin 5% + Copper Oxychloride 45% WP | ● | Fluxapyroxad 7.5% + Difenoconazole 5% SC | ● |
| Hexaconazole 5 % SC | ● | Pyraclostrobin 33.3 % + Fluxapyroxad 16.7 % SC | ● |
| Tebuconazole 25% EC | ● | Dimethomorph 50% WP | ● |
| Mancozeb 75% WP | ● | Cymoxanil 8% + Mancozeb 64% WP | ● |
| Fluxapyroxad 25% + Pyraclostrobin 25% SC | ● | Azoxystrobin 8.3% + Mancozeb 66.7% WDG | ● |
| Tebuconazole 50% + Trifloxystrobin 25% WG | ● | Dimethomorph 12% + Pyraclostrobin 6.7% WG | ● |
| Metiram 70% WG | ● | Fluopicolide 5.56 + Propamocarb Hydrochloride 55.6% SC | ● |
| Metalaxyl 4% + Mancozeb 64% WP | ● | Iprovalicarb 5.5% + Propineb 61.25% WP | ● |
| Carbendazim 12% + Mancozeb 63% WP | ● | Fenamidone 10% + Mancozeb 50% WG | ● |
| Difenoconazole 25% EC | ● | Ametocetradin 27% + Dimethomorph 20.27% SC | ● |
| Fosetyl Al 80% WP | ● | Metalaxyl-M 3.3% + Chlorothalonil 33.1% SC | ● |
| Propineb 70% WP | ● | Chlorothalonil 75 % WP | ● |
| Tetraconazole 3.8% EW | ● | Mandipropamid 25% SC | ● |
| Captan 70% + Hexaconazole 5% WP | ● | Sulphur 80% WDG | ● |
| Thiophanate Methyl 70% WP | ● | Azoxystrobin 23% SC | ● |
| Azoxystrobin 11% + Tebuconazole 18.3% SC | ● | Zineb 75% WP | ● |
| Kresoxim-Methyl 44.3% SC | ● | | |

PGR AND MICRO-NUTRIENTS

| COMPATIBILITY WITH EFFICON® | |
|--|---|
| 00:52:34 | ● |
| 13:00:45 | ● |
| 19:19:19 | ● |
| Biovita-Ascophyllum nodosum | ● |
| Fantac-Mixture of Amino acids and vitamins | ● |
| Librel Zinc | ● |
| Isabion-Amino acids | ● |
| Librel TMX2 | ● |
| Lihocin | ● |
| Biozyme-Seaweed | ● |



● Compatible with Efficon

○ Noticed marginal crop response and recovered within 5 to 7 days. (Yellow patches & slight shrinking of leaves, noticed only in Brinjal & Cucumber)

○ Not Recommended: Shows Crop Response like Bleaching & Chlorosis on leaves, usually recovered in two-to-three-week time. Noticed only in Cotton.

BASF

We create chemistry